

## Addition of Risk Factors of Coronary Heart Disease in Diagnostic Value of Treadmill Score for Detecting Complexity of Coronary Arterial Lesions

Mariyetty K.S. Nasution\*, Abdullah A. Siregar, Harris Hasan, Zulfikri Mukhtar, Abdul H. Raynaldo, Nizam Akbar

Department of Cardiology and Vascular Medicine, Faculty of Medicine University of Sumatera Utara, Adam Malik Hospital, Medan, Indonesia

\*Corresponding author :

Mariyetty Kemuning Sari Nasution, MD, - email: kemuning29@gmail.com

Department of Cardiology and Vascular Medicine, University of Sumatera Utara, Adam Malik Hospital Medan, Indonesia, JalanBunga Lau, Medan 55281, Indonesia

Manuscript submitted: August 20, 2018; Revised and accepted: October 3, 2018

### ABSTRACT

**Background:** Coronary heart disease (CHD) is still the leading cause of death in the world. There are various risk factors for atherosclerosis leading to CHD. Duke Treadmill Score (DTS) is known to demonstrate prognostic stratification and has a diagnostic value in predicting the number of coronary arteries involved in patient populations with ischemic heart disease. However, DTS does not describe the role of risk factors for coronary heart disease to the complexity of coronary artery lesions. This study aims to add risk factors for coronary heart disease on DTS to detect the complexity of coronary artery lesions with stable angina pectoris.

**Methods:** This study was a cross-sectional study in stable angina pectoris patient who comes to Haji Adam Malik Hospital Medan from January 2017 until February 2018. Patients who have done treadmill test and coronary angiography, and fulfill inclusion and exclusion criteria are included in the study. ECG examination and recording of risk factors for coronary heart disease were done. DTS assessment was performed based on a treadmill test and Syntax score based on coronary angiography results. Diagnostic tests were performed to assess the sensitivity and specificity of the addition of CHD risk factors to detect the complexity of coronary artery lesions.

**Results:** Of the 76 people with stable angina pectoris, 55 people were found with low SYNTAX and 21 people with high Syntax. DTS is divided into 3 groups: mild ( $> -10$ ), moderate ( $-10$  to  $-13.5$ ), and severe ( $\leq -13.6$ ) based on the cut off of the ROC curve. Risk factors for CHD are divided into 3 groups, mild ( $\leq 3$  CHD risk factors), moderate (4-6 CHD risk factors), and severe (7 CHD risk factors) based on the cut off of the ROC curve, then assessed the relationship with Syntax which has been divided into 2 groups, low Syntax, and high Syntax. Diagnostic test shows the addition of risk factors of CHD to DTS to detect the complexity of coronary artery lesions have greater sensitivity and specificity than DTS without the addition of risk factors of CHD, 95%, and 89%.

**Conclusion:** The addition of risk factors for coronary heart disease on DTS can detect the complexity of coronary artery lesions.

**Keywords:** Duke Treadmill Score; risk factors; complexity of coronary arterial lesions; Syntax.

### INTISARI

**Latar Belakang:** Penyakit jantung koroner (PJK) masih merupakan penyebab kematian utama di dunia. Terdapat berbagai factor resiko terjadinya aterosklerosis yang mendorong terjadinya PJK. Duke Treadmill Score (DTS) diketahui dapat menunjukkan stratifikasi prognostic dan memiliki nilai diagnostic dalam memprediksi jumlah arteri koroner yang terlibat pada populasi penderita dengan penyakit jantung iskemik. Namun, DTS tidak menggambarkan peran factor resiko penyakit jantung koroner terhadap kompleksitas lesi arteri koroner. Penelitian ini

bertujuan untuk menambahkan factor resiko penyakit jantung koroner pada nilai DTS untuk mendeteksi kompleksitas lesi arteri koroner penderita angina pectoris stabil.

**Metode:** Penelitian ini merupakan penelitian potong lintang terhadap penderita angina pectoris stabil yang datang ke poliklinik Rumah Sakit Haji Adam Malik Medan sejak Januari 2017 sampai Februari 2018. Penderita yang telah dilakukan uji latih jantung dan angiografi koroner, dan memenuhi criteria inklusi dan eksklusi dimasukkan ke dalam penelitian. Dilakukan pemeriksaan EKG dan pencatatan factor resiko penyakit jantung koroner. Penilaian DTS dilakukan berdasarkan uji latih jantung dan skor Syntax berdasarkan hasil angiografi koroner. Dilakukan uji diagnostik untuk menilai sensitivitas dan spesifisitas penambahan factor resiko PJK untuk mendeteksi kompleksitas lesi arteri koroner

**Hasil:** Dari 76 orang penderita angina pectoris stabil, didapatkan 55 orang dijumpai dengan Syntax rendah dan 21 orang dengan Syntax tinggi. DTS dibagi dalam 3 kelompok yaitu ringan ( $>-10$ ), sedang ( $-10$  sampai dengan  $-13,5$ ), dan berat ( $\leq-13,6$ ) berdasarkan titik potong kurva ROC. Faktor resiko PJK dibagi ke dalam 3 kelompok, yaitu ringan ( $\leq 3$  faktor resiko), sedang (4-6 faktor resiko), dan berat (7 faktor resiko) berdasarkan titik potong kurva ROC, kemudian dinilai hubungannya dengan Syntax yang telah dibagi menjadi 2 kelompok, yaitu Syntax rendah dan Syntax tinggi. Hasil uji diagnostic menunjukkan penambahan factor resiko pada nilai DTS untuk mendeteksi kompleksitas lesi arteri koroner memiliki sensitivitas dan spesifisitas yang lebih besar dibandingkan DTS tanpa penambahan factor resiko PJK, yaitu 95% dan 89%

**Kesimpulan:** Berdasarkan hasil analisis pada penelitian ini, dapat disimpulkan bahwa penambahan faktor resiko penyakit jantung koroner pada nilai diagnostik DTS dapat mendeteksi kompleksitas lesi arteri koroner.

## INTRODUCTION

Coronary heart disease (CHD) is a condition where there is an imbalance between oxygen supply and the needs that cause hypoxic conditions in the myocardium and accumulation of metabolic waste substances that are generally caused by the process of atherosclerosis in the coronary arteries.<sup>1</sup> CHD is the leading cause of death in the world. A report from the World Health Organization (WHO) in 2013, 17.5 million people dies each year from cardiovascular disease, about 31% of the total deaths worldwide.<sup>2</sup> There are various risk factors for atherosclerosis that encourage CHD, both modifiable risk factors such as diabetes mellitus, hypertension, smoking, lack of activity and dyslipidemia, or those that cannot be modified such as age, sex, and family history.<sup>3</sup> The most common clinical presentation of CHD is stable angina pectoris followed by acute coronary syndrome and then sudden death. Treadmill test is the most widely used test to conduct an initial evaluation of patients with angina pectoris.<sup>4</sup> Prognostic value of treadmill test based on exercise duration, ST segment deviation (depression or elevation), and the presence and severity of angina that arises during exercise. This value is known as Duke Treadmill Score (DTS) which shows prognostic stratification

in a population of patients with a history of miocard infarction and with typical chest pain.<sup>5,6</sup>

There are several value systems used to describe the complexity of coronary artery lesions. The commonly used value system is Syntax (Synergy between percutaneous coronary intervention with Taxus and cardiac surgery). Syntax can objectively determine the upper limit of suitability for revascularization with percutaneous intervention methods or revascularization methods with coronary bypass surgery.<sup>7</sup> Duke Treadmill Score (DTS) has been shown to have a strong correlation with the complexity of coronary artery lesions assessed by Syntax.<sup>8,9</sup> However, DTS does not describe the role of risk factors for coronary heart disease in the complexity of coronary artery lesions, there is no data regarding the addition of risk factors for CHD in DTS diagnostic values in predicting the complexity of coronary artery lesions assessed based on Syntax.

## METHODS

This study is a diagnostic study with cross-sectional design. This study uses Treadmill test data, history of risk factors for coronary heart disease, and coronary angiography at Haji Adam Malik Hospital (RSHAM) from January 2017 to February 2018. Patients who present with stable

angina pectoris CCS II-III and sinus rhythm, can perform treadmill test and no contraindication to treadmill test and coronary angiography is included in the study. Patients who cannot perform treadmill test, refuse to do coronary angiography, chronic kidney disease, a history of severe valvular disease, congenital heart disease, heart failure, acute coronary syndrome, coronary heart disease, myocardial infarction, or previously revascularization is excluded in this research. The independent variable in this study is DTS and CHD risk factors including diabetes mellitus, hypertension, dyslipidemia, BMI, smoking, gender, and age. CHD risk factors will be divided into mild risk factors ( $\leq 3$  CHD risk factors), moderate (4-6 CHD risk factors), severe (7 CHD risk factors) based on the Receiver Operating Characteristic (ROC) cut off point. The dependent variable in this study is the complexity of coronary artery lesions which are assessed by Syntax score which is then divided into low Syntax ( $\leq 22$ ) and high Syntax ( $> 22$ ).

Treadmill test in this study used Bruce protocol. The DTS is obtained based on the calculation of exercise duration (in minutes) - 5x (deepest ST segment deviation in mm) -4x (angina score).<sup>10</sup> The duration of the exercise used in the calculation is based on the duration of the exercise in minutes on the Bruce protocol.

#### Statistical Analysis

Statistical data processing and analysis are using statistical computer devices. Categorical variables are presented with the number or frequency (n) and percentage (%). Numerical variables are presented with mean values with standard deviations for data that are normally distributed. Test the normality of numerical variables in all study subjects using the Kolmogorov-Smirnov test with  $n > 50$ . Comparison between the two groups in categorical independent variables and categorical dependent variables using Chi-Square test ( Fisher test). Comparison between the two groups on numerical independent variables and category dependent variables is done by independent T test ( Mann Whitney test). We used receiver operating characteristic analysis to determine the optimal cut off point of DTS. Using Chi-squared table, we obtained sensitivity, specificity, positive

predictive value (PPV) and negative predictive value (NPV). The association of the DTS test results on the complexity of coronary artery lesions will be compared with the association of the DTS test results with additional risk factors for CHD in the form of sensitivity and specificity.<sup>11</sup> All statistical analyses were performed using SPSS, version 22.0 and p value  $< 0.05$  was considered as statistically significant.

## RESULTS

This study took place from January 2017 to February 2018. The total number of study who met the inclusion and exclusion criteria were 76 subjects consisting of 21 people with high Syntax and 55 people with low Syntax. Male gender consisted of 19 people (90.5%) in the group with high Syntax and 37 people (67.3%) in the low Syntax. The mean age of 54.43 years was in the group with high Syntax and 53.36 years in the low Syntax. A bivariate analysis was carried out on the factors that influenced Syntax in both groups. In bivariate analysis, CHD risk factors such as hypertension, dyslipidemia, BMI, sex, age, and diabetes mellitus did not have a significant difference between groups with high Syntax and low Syntax.

In addition, there were no significant differences between medication history, and laboratory values such as HbA1C, fasting blood glucose, add random blood glucose, 2 hours postprandial blood glucose, HDL, LDL and triglycerides. CHD risk factors especially dyslipidemia and mean DTS have statistically significant differences in the two groups of Syntax. In the high Syntax group there were 15 (71.4%) subjects with dyslipidemia and 21 (38.2%) subjects with dyslipidemia have low Syntax ( $p = 0.011$ ). The mean DTS in the high Syntax group was  $-18.63 \pm 8.85$  and the low Syntax was  $-9.21 \pm 3.41$  ( $p = 0.000$ ) (Table.1).

Syntax assessment was carried out by two cardiologist consultants who first performed the kappa interobserver test to find the suitability of the interpretation. From the kappa test, it was obtained a value of 0.773 with  $p < 0.001$ , meaning that there was a good match between the first and second observations.

Analysis with the ROC curve is done to find out the diagnostic value of DTS to predict high syntax. The area under the

curve (AUC) was 90.1% (95% CI 0.804-0.999,  $p = 0.000$ ) (figure 1). After that, the determination of the DTS is based on the ROC curve. The limit value is determined based on the point of intersection between the optimal sensitivity and specificity to obtain high Syntax. A DTS-13.6 is considered to have optimal diagnostic value.

Further analysis is carried out to determine the diagnostic value of DTS with additional risk factors for CHD to detect the complexity of coronary artery lesions. DTS with additional risk factors for CHD is divided into 2 groups, The low risk which is

classified as mild DTS accompanied by mild, moderate or severe CHD risk factors, and the high risk which is classified as moderate or severe DTS accompanied by mild, moderate or severe CHD risk factors. Thus 20 people classified as high risk have a high Syntax, while 49 people with low risk have a low Syntax. The calculation of diagnostic values is carried out based on Chi-square table and it is obtained a sensitivity of 95%, specificity of 89%, negative predictive value of 98%, and positive predictive value of 77% ( $p < 0.001$ ) (table 2).

**Table 1. Demographic and clinical characteristics based on Syntax Score**

Characteristics	High Syntax (n=21)	Low Syntax (n=55)	P value
Gender			
Men, n (%)	19(90.5)	37(67.3)	0.040
Women, n (%)	2(9.5)	18(32.7)	
Age (mean±SD), years	54.43±6.80	53.36±8.15	0.567
Heart Rate(mean±SD),x/min	80.19±8.98	78.36±8.59	0.428
DBP (mean±SD), mmHg	82.38±13.00	85.09±11.99	0.392
SBP (mean±SD), mmHg	134.76±10.30	138.51±13.05	0.241
BMI (mean±SD), kg/m <sup>2</sup>	25.09±2.70	26.13±3.38	0.172
<b>Risk factors CHD</b>			
Diabetes Mellitus, n (%)	8 (38.1)	15(27.3)	0.358
Hypertension, n (%)	16(76.2)	45(81.8)	0.581
Smoking, n (%)	17 (81.0)	35(63.6)	0.043
Dyslipidemia, n (%)	13(61.9)	18(32.7)	0.021
Age, n (%)	18 (85.7)	44 (80.0)	0.745
Long time for Diabetes Mellitus, (mean±SD)	1.48±3.07	1.04±2.30	0.557
Duke Treadmill Score, (mean±SD)	-19.95±8.28	-10.56±4.60	0.000
<b>Laboratory</b>			
HbA1C (mean±SD), %	7.28±1.74	7.30±6.52	0.992
Fasting blood glucose (mean ±SD), mg/dL	133.85±44.19	114.72±26.64	0.075
Random blood glucose (mean±SD), mg/dL	162.42±116.4	122.72±49.36	0.145
2 h p.pblood Glucose (mean±SD), mg/dL	181.90±67.8	180.12±58.11	0.916
Triglycerides (mean±SD), mg/dL	135.66±48.55	134.40±47.6	0.918
Total Cholesterol(mean±SD),mg/dL	188.23±50.19	182.12±38.72	0.618
HDL (mean±SD), mg/dL	40.85±10.01	44.54±8.11	0.141
LDL (mean±SD), mg/dL	126.57±51.04	117.76±33.90	0.385
<b>History of Medication (n,%)</b>			
Diuretic,n (%)	1(4.8)	1(1.8)	0.479
ACE Inhibitors/ARB, n(%)	15 (71.4)	45 (81.8)	0.320
Nitrate, n(%)	19 (90.5)	46 (83.6)	0.449




Figure 1.The ROC curve of DTS value to assess Syntax score

**Table 2. Chi-squared test based on DTS value with additional risk factors for CHD**

DTS with additional risk factors	High Syntax	Low Syntax	Total	Sensitivity	Specificity	NPV	PPV	P value
High Risk	20	6	26	95%	89%	98%	77%	0.000
Low Risk	1	49	50					
Total	21	55	76					

**DISCUSSION**

Of the total subjects in this study, men were the highest subjects compared to women, 59 subjects (77.6%) compared to 17 subjects (22.4%) and had a higher percentage of Syntax than women, 19 subjects (90.5%) compared to 2 subjects (9.5%). Based on a population study conducted by Cheng *et al.*<sup>12</sup>It was found that men with typical angina had a greater risk factor for coronary heart disease with severe lesions than women (19% versus 11% with  $p < 0.001$ ). The average age of this study is 60 years, according to previous studies with an average age ranging from 50-60 years.<sup>13</sup> At the results of risk factors such as hypertension, diabetes melitus, smoking, BMI, age, and gender, are more common in high Syntax compared to low Syntax, but not statistically significant. Risk factors for dyslipidemia are different and meaningful to Syntax ( $p < 0.05$ ). This is consistent with data from the Framingham study and other cohort studies which show

that there is a risk of ischemic heart disease in dyslipidemia in the form of an increase in total cholesterol. Coronary risk occurs as much as 2 times as in someone with a cholesterol level of 240 mg / gL compared to someone with a cholesterol level of 200 mg / dL).<sup>14</sup>From the results of the bivariate analysis the group of subjects with high Syntax had lower DTS mean scores than the low Syntax group, and this was statistically significant. The mean DTS in the high Syntax group was  $-18.63 \pm 8.85$  and the low Syntax was  $-9.21 \pm 3.41$  ( $p < 0.001$ ). These results are consistent with previous studies which stated that there was a significant negative correlation between DTS and Syntax ( $r = -0.9$ ),  $p < 0.001$ ), where subjects included in the high risk DTS had a higher mean Syntax than DTS moderate risk and low risk.

From the DTS diagnostic value based on the ROC curve, the area under the curve is 90.1% ( $p < 0.001$ ) and based on the area under this curve the diagnostic value of DTS is very good.<sup>8</sup> Further analysis

is carried out to determine the DTS limit value which has an optimal diagnostic value to predict Syntax values. DTS-13.6 is considered to have optimal diagnostic value because it has a balanced sensitivity and specificity. Based on these basic values, DTS is classified into two groups, DTS $\leq$ -13.6 and DTS $>$  -13.6. Calculation of diagnostic values is done based on Chi-squared table and obtained a sensitivity of 90%, specificity of 61%, negative predictive value of 91.6%, and positive predictive value of 60%. The results of this study have better sensitivity and specificity than studies conducted by Gunaydin *et al.*<sup>9</sup> who assessed the DTS relationship with the severity of coronary artery disease assessed by Syntax and obtained AUC of 0.83 (IK 0.7-0.88) with sensitivity of 74 % and specificity of 73%.<sup>4</sup> The diagnostic value of DTS sensitivity obtained in this study is also better than the diagnostic value based on study conducted by Tamargo *et al.*<sup>15</sup>, where high risk DTS (DTS  $<$ -11) has a sensitivity of 30.8%, specificity of 90.5%, for detecting coronary lesions severity. However, the assessment of coronary artery lesions in the Tamargo study was based on the number and type of coronary arteries involved, whereas in this study were based on Syntax.

Although the diagnostic value of DTS is generally effective for stratifying patients with stable angina pectoris into the appropriate diagnostic and prognostic categories, the addition of other prognostic covariates such as risk factors for coronary heart disease is expected to increase the accuracy of their predictions.

Further analysis was carried out to assess DTS which has added risk factors for CHD to detect the complexity of coronary artery lesions. DTS with risk factors for CHD is divided into 2 groups, high and low risk. Calculation of the diagnostic value is based on Chi-squared table and obtained a sensitivity of 95%, specificity of 89%, negative predictive value of 98%, and positive predictive value of 77%. The results of this study support the study hypothesis that the DTS values with risk factors for CHD have good diagnostics to detect the complexity of coronary artery lesions which in this study were defined as coronary lesions that have high Syntax . DTS with risk factors for CHD has better sensitivity and specificity than DTS without the addition of CHD risk factors.

Adding risk factors for CHD to DTS values can help in stratifying stable angina pectoris with a high risk group, that will benefit from revascularization and its clinical application is also easy and can be performed on all health services because the variables used are simple and can be done on all health care systems. Limitations of this study include the absence of a time limit between the treadmill test and coronary angiography at the time of selection of subjects that can be related to the progression of coronary lesion severity. In addition, the percentage of subjects with low female sex caused a low representation of female sex in this study. This study only assessed the number of CHD risk factors and their relationship with Syntax, but did not assess risk factors for CHD as a predictor of Syntax. Prospective studies with larger samples are needed to confirm the efficacy of DTS that has added risk factors for CHD to detect the complexity of coronary artery lesions. The addition of CHD risk factors to DTS values to detect the complexity of coronary artery lesions is very simple and easy to do, so we need system validation.

## CONCLUSION

Among all CHD risk factors in this study, only dyslipidemia was different and statistically significant against Syntax ( $p < 0.05$ ). DTS which has added risk factors for CHD has a higher sensitivity and specificity, 95% and 85% compared to DTS without the addition of CHD risk factors, 90% and 61% to detect the complexity of coronary artery lesions.

## REFERENCES

1. Wilder J., Sabatine M.S., Lilly L.S. 2016. Ischemic Heart Disease Chapter 6. In: Lilly, Leonard S. Pathophysiology of heart disease: a collaborative project of medical students and faculty. 6th ed. Baltimore: Lippincott Williams & Wilkins, pp.134-161.
2. Haris H., Elvrida D. 2016. Angina Pectoris Stabil (APS). In: Kasiman (eds) Nyeri dada dari A sampai Z. Medan: Pustaka Bangsa, pp. 10-40.
3. Lilly L.S. 2012. Pathophysiology of heart disease: a collaborative project of medical students and faculty. 6th ed.

- Baltimore: Lippincott Williams & Wilkins.
4. Tardif C.J. 2010. Coronary artery disease in 2010. *Eur Heart J Supplement*, 12 (Supplement C), C2–C10.
  5. Gibbons R.J., Balady G.J., Bricker J.T., Chaitman B.R., Fletcher G.F., Froelicher V.F., et al. ACC/AHA 2002 Guideline update for exercise testing. American College of Cardiology Foundation and American Heart Association. *J Am CollCardiol*, 40:1531-1540.
  6. Mark D.B., Hlatky M.A., Harrell F.E.Jr., Lee K.L., Califf R.M., Pryor D.B. 1987. Exercise treadmill score for predicting prognosis in coronary artery disease. *Ann Intern Med*, 106:793-800.
  7. Mohr F.W., Morice M.C., Kappetein A.P., Feldman T.E., Stähle E., Colombo A., et al. 2013. Coronary artery bypass graft surgery versus percutaneous coronary intervention in patients with three-vessel disease and left main coronary disease: 5-years follow up of the randomized, clinical SYNTAX Trial. *Lancet*, 381:629-638.
  8. Acar Z., Korkmaz L., Agac M.T., Erkan H., Dursun I., Kalaycioglu E., et al. 2012. Relationship between Duke treadmill score and coronary artery lesion complexity. *Clin Invest Med*, 35:E365-369.
  9. Günaydın Z.Y., Bektaş O., Gürel Y.E., Karagöz A., Kaya A., Kırış T., et al. 2016. The value of the Duke treadmill score in predicting the presence and severity of coronary artery disease. *Kardiol Pol*, 74:127-134.
  10. Perhimpunan Dokter Spesialis Kardiologi Indonesia. 2016. Pedoman Uji Latih Jantung: Prosedur dan Interpretasi. Jakarta: PERKI.
  11. Dahlan M.S. 2009. Penelitian Diagnostik. Edisi ketiga. Jakarta: Penerbit Salemba Medika.
  12. Cheng V.Y., Berman D.S., Rozanski A., Dunning A.M., Achenbach S., Al-Mallah M., et al. 2011. Performance of the traditional age, sex, and angina typicality-based approach for estimating pretest probability of angiographically significant coronary artery disease in patients undergoing coronary computed tomographic angiography. *Circulation*, 124:2423-2432.
  13. Mayasari N.M.E, Arso I.A., Maharani E. 2016. Nilai diagnostik Duke treadmill score untuk mendeteksi keparahan lesi arteri koroner pada pasien yang terduga penyakit jantung koroner stabil. *J Kardiol Indones*, 37:65-74
  14. Shahawy S., Libby P. 2016. Atherosclerotic. In: Lilly, Leonard S. Pathophysiology of heart disease: a collaborative project of medical students and faculty. 6th ed. Baltimore: Lippincott Williams & Wilkins, pp. 112-133.
  15. Alvarez-Tamargo J.A., Simarro Martín-Ambrosio E., Romero-Tarín E., Albadalejo-Salinas V.J., de la Hera J.M., Martín-Fernández M., et al. 2005. Angiographic evaluation of high risk treadmill score in patient with stable angina according to sex, age, or use of drugs with negative chronotropic effect. *Rev Esp Cardiol*, 59:448-457.