

**VIABILITY TEST OF STORED CORN SEEDS BASED ON THE RELATIONSHIP
BETWEEN CONCENTRATION OF INORGANIC EXUDATES AND ELECTRICAL
CONDUCTIVITY OF SOAKING WATER**

**UJI VIABILITAS BENIH JAGUNG
BERDASARKAN HUBUNGAN KONSENTRASI EKSUDAT ANORGANIK
DENGAN DAYA HANTAR LISTRIK AIR RENDAMAN**

T. S. Endang¹, P. Yudono², and D. Indradewa²

INTISARI

Penelitian ini dilakukan untuk mengetahui hubungan konsentrasi eksudat dengan daya hantar listrik air rendaman benih jagung setelah disimpan dalam rangka uji mutu benihnya. Rancangan Acak Lengkap digunakan untuk perlakuan periode penyimpanan 0, 1, 2, 3, 4 bulan dalam suhu kamar. Ulangan dilakukan tiga kali. Pengamatan dilakukan terhadap konsentrasi eksudat anorganik (nitrogen, fosfor, kalium, kalsium, dan magnesium), daya hantar listrik, daya tumbuh benih, serta bobot basah dan kering bibit.

Hasil penelitian menunjukkan bahwa dengan semakin lamanya penyimpanan, konsentrasi eksudat anorganik meningkat, daya hantar listrik meningkat, tetapi daya tumbuh benih dan bobot bibit menurun. Konsentrasi eksudat memiliki korelasi positif yang nyata terhadap daya hantar listrik dan memiliki korelasi negatif yang nyata terhadap daya tumbuh benih dan bobot bibitnya. Hasil ini memungkinkan penggunaan daya hantar listrik sebagai indikator viabilitas benih dan alternatif cara uji daya tumbuh benih jagung.

Kata kunci: elektrolit, eksudat anorganik, daya hantar listrik, viabilitas benih, jagung.

ABSTRACT

It has been known that deteriorate seeds exudes much ions if soaked in water. This experiment was aimed to assess stored corn seeds viability based on concentration of inorganic exudates and on electrical conductivity (EC) of soaking water of the seeds they generate.

Seeds of zero, one, two, three, and four months in storage were soaked in water. The soaking water then was observed for its nitrogen, phosphorous, potassium, calcium, and magnesium concentration, as well as their respective electrical conductivity. Simultaneously, seeds germination percentage and seedling weight was also observed.

The results indicated that concentration of inorganic exudates and the EC of the soaking water had close and negative relationship with germination percentage and seedling weight along the storage periods examined. Based on this finding, the EC could be used as an alternative method for viability test of corn seeds.

Keywords : electrolite, inorganic exudate, electrical conductivity (EC), seed viability, corn.

¹ Alumni Fakultas Pertanian Universitas Gadjah Mada

² Staf pengajar Fakultas Pertanian Universitas Gadjah Mada

INTRODUCTION

Quality of seeds consists of genetical, physiological, and physical aspects. Those aspects are the key factors of seed germination and seedling growth. Seed viability measures the physiological aspect of seed quality, and manifests in ultimate metabolic processes, such as seed germination, vigor, and seedling growth (Sadjad, 1975).

Seed deterioration, as an indication of seed viability declination, is usually affected by unfavorable condition of storage room, length of storage, and internal condition of the seed itself. It was characterized by low germination capacity, vigor, and growth of seedling. The process is irreversible and starts soon after seeds reach physiological maturity (Abdul Baki and Anderson, 1972). Suseno (1975) mentioned that germination capacity and seedling vigor are widely used as measures of seed deterioration.

Bewley and Black (1978) stated that the degradation of cell membrane integrity in seed deterioration resulting in the control function loss towards membranes permeability. This condition affects the outflow of exudates from cells. The concentration of exudates (organic and inorganic) will be found in seeds soaking water, thus it relates with the degree of seed deterioration. The exudates mainly contain inorganic cations and organic compounds that have electrical conductivity (EC) (Copeland, 1976). If the amount of exudates has any relationship with the EC, we may measure the seed deterioration using this feature.

This experiment was conducted to obtain proofs on the relationship of electrical conductivity with the exudates as inorganic cations concentration, so that can be used as an indirect measurement of seed deterioration

MATERIALS AND METHODS

Newly harvested good quality corn seeds were obtained from Cereals Seed Production Station Soropadan, Central Java. Then the seeds were stored for one, two, three and four months under the condition of room temperature, placed in polyethylene bags. The moisture content of the seeds at the start of storage was 12.0 percent.

The design used in this experiment was Randomized Completely Design (CRD), single factor (periods of storage) three replications. Observations were done on the germination percentage, the electrical conductivity of the seeds soaking water, the concentration of inorganic exudates as Nitrogen, Phosphorous, Potassium, Calcium, and Magnesium in the seeds soaking water, and the weight of seedlings.

The data obtained were analysed statistically using varianc analysis of and the Duncan's New Multiple Range Test (Gomez and Gomez, 1976).

RESULTS AND DISCUSSION

The obtained data and the results of the analysis of variance on the germination percentage, fresh and dry weight of seedlings, electrical conductivity and the concentrations of Nitrogen,

Table 1. The Mean values of germination percentage (%), fresh weight of seedling (g), dry weight of seedlings (g), electrical conductivity (mmhos), concentration of inorganic exudates (Nitrogen, Phosphorus, Kalium, Calcium, and Magnesium (ppm)) in seeds soaking water, observer after storage.

Variable	Periods of Storage (Month)				
	0	1	2	3	4
Germination percentage	100 ^a	97 ^a	94 ^a	91 ^{bc}	87 ^c
Fresh weight	8.94 ^a	8.46 ^a	8.00 ^{ab}	6.79 ^b	6.25 ^c
Dry weight	4.47 ^a	4.37 ^a	3.97 ^a	2.66 ^b	2.38 ^c
Electrical conductivity	0.30 ^a	0.32 ^b	0.33 ^{bc}	0.35 ^c	0.64 ^d
Nitrogen	5.6 ^a	12.1 ^b	13.3 ^b	14.7 ^b	15.4 ^b
Phosphorous	1.64 ^a	1.92 ^b	3.33 ^b	4.08 ^b	7.18 ^c
Potassium	3.23 ^a	3.65 ^b	9.43 ^b	10.1 ^c	14.99 ^d
Calcium	0.06 ^a	0.07 ^b	0.13 ^b	0.31 ^b	1.05 ^c
Magnesium	0.18 ^a	0.20 ^{ab}	0.24 ^b	0.31 ^b	0.43 ^c

Notes: Mean values followed by the same letter in the same row are not significantly different (DMRT 5%).

Table 2. The correlations and regressions between the periods of storage (X) and the germination percentage, EC, fresh weight, dry weight of seedlings, concentration of Nitrogen, Phosphorus, Potassium, Calcium and Magnesium in seeds soaking water.

Dependent Variable	Correlation Coefficient	Regression Equation
Germination percentage	-0.999 *	$Y = -7.1x + 100.0$
Fresh weight	-0.985 *	$Y = -7.9x + 9.19$
Dry weight	-0.950 *	$Y = -0.58x + 4.71$
Electrical conductivity	-0.790 *	$Y = -0.071x + 0.25$
Concentration of Nitrogen	0.897 *	$Y = 0.22x + 7.78$
Concentration of Phosphorus	0.945 *	$Y = 1.32x + 0.982$
Concentration of Kalium	0.965 *	$Y = 3.0x + 2.29$
Concentration of Calcium	0.840 *	$Y = 9.22x + -0.12$
Concentration of Magnesium	0.950 *	$Y = 0.06x + 0.02$

Note : Significance at 5% level.

Table 3. The correlations and the regressions between the EC and the germination percentages, germination rate, and field emergences.

Variables	Correlation Coefficient	Regressions Equation
Germination percentages	-0.674 *	$Y = -0.023x + 2.59$
Germination rate	-0.810 *	$Y = -0.028x + 1.139$
Field emergences	-0.795 *	$Y = -0.029x + 3.154$

Note : Significance at 5% level.

Phosphorous, Potassium, Calcium, and Magnesium were presented in Table 1, and analysis correlation and regression results were presented in Table 2.

It was shown that the storage caused deterioration of seeds, denoted by increase of the concentration of the inorganic exudates in the seeds soaking water. The increment of these elements causing higher electrical conductivity, since the elements are electrolytes. The leakages were already significant starting from the second month of storage, and steadily increased up to the termination of the storage. The disintegration of the membranes were followed by the decrease of the viability of the seeds, manifested in seedling germination percentage, dry and fresh weight.

The results proved that germination percentage, fresh and dry weight of seedlings indicated significant positive correlations, while the other variables showed significant negative correlations.

Table 3 shows the correlation and regressions between the EC and the germination percentages,

germination rate and the field emergences. It indicated that the EC of the seeds soaking water were significantly correlated to the lower germination and the field emergences of the corn seeds after storages.

Further analysis found that the correlations between EC and the concentration of inorganic exudates (Nitrogen, Phosphorus, Potassium, Calcium and Magnesium) were : $r = 0.5719^*$; $r = 0.9390^*$; $r = 0.8259^*$; $r = 0.9904^*$; and $r = 0.9388^*$ respectively. The elements, as electrolytes, gave the contributions to the EC of seeds soaking water of the aged corn seeds (Skoog and West, 1969).

CONCLUSION

The concentrations of the inorganic exudates increased with the longer the period of storages indicated the progress of the seed deterioration. The EC highly correlated with concentration of inorganic exudates, as germination percentages and the field emergences. The results also indi-

cated that the EC could be used as an alternative method for corn seeds viability tests.

LITERATURE CITED

- Abdul Baki, A. A. and J. D. Anderson. 1972. Physiological and Biochemical Deterioration of Seeds. In: T. T. Kozlowski. (ed.). *Seed Biology II*. New York: Academic Press. pp 283 - 309.
- Bewley, J. D. and M. Black. 1978. *Physiology and Biochemistry of Seeds I*. Springer-Verlag, Berlin - Heidelberg. New York.
- Copeland, L. O. 1976. *Principles of Seed Science and Technology*. Burgess Publishing. Minneapolis.
- Gomez, K. A. and A. A. Gomez. 1976. *Statistical Procedures for Agricultural Research*. Los Banos.
- Sadjad, S. 1975. Teknologi Benih dan Masalah-Masalahnya. In Sadjad (ed.). *Dasar-Dasar Teknologi Benih*. IPB Bogor. pp: 1-11.
- Skoog, D. A. and D. M. West. 1969. *Fundamental of Analytical Chemistry*. Holt Rinehart and Winston Inc. New York.
- Suseno, H. 1975. Fisiologi dan Biokimia Kemunduran Benih. In: S.Sadjad (ed.) *Dasar-Dasar Teknologi Benih*. IPB Bogor. pp : 98 - 126.